



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
General Certificate of Education
Advanced Subsidiary Level and Advanced Level

Mathematics (9709)

Paper 3: Pure Mathematics (P3)

2020-2021



UNIVERSITY *of* CAMBRIDGE
International Examinations



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



MATHEMATICS

9709/32

Paper 3 Pure Mathematics 3

February/March 2020

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Blank pages are indicated.

1 (a) Sketch the graph of $y = |x - 2|$.

[1]

(b) Solve the inequality $|x - 2| < 3x - 4$.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 2 Solve the equation $\ln 3 + \ln(2x + 5) = 2 \ln(x + 2)$. Give your answer in a simplified exact form. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 3 (a) By sketching a suitable pair of graphs, show that the equation $\sec x = 2 - \frac{1}{2}x$ has exactly one root in the interval $0 \leq x < \frac{1}{2}\pi$. [2]

.....

.....

- (b) Verify by calculation that this root lies between 0.8 and 1. [2]

.....

.....

.....

.....

.....

- (c) Use the iterative formula $x_{n+1} = \cos^{-1}\left(\frac{2}{4-x_n}\right)$ to determine the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

4 Find $\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} x \sec^2 x \, dx$. Give your answer in a simplified exact form. [7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence solve the equation $\frac{\cos 3x}{\sin x} + \frac{\sin 3x}{\cos x} = 4$, for $0 < x < \pi$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) State what happens to the value of y as x tends to infinity. [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

7 The equation of a curve is $x^3 + 3xy^2 - y^3 = 5$.

(a) Show that $\frac{dy}{dx} = \frac{x^2 + y^2}{y^2 - 2xy}$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



In the diagram, $OABCDEFG$ is a cuboid in which $OA = 2$ units, $OC = 3$ units and $OD = 2$ units. Unit vectors \mathbf{i} , \mathbf{j} and \mathbf{k} are parallel to OA , OC and OD respectively. The point M on AB is such that $MB = 2AM$. The midpoint of FG is N .

- (a) Express the vectors \overrightarrow{OM} and \overrightarrow{MN} in terms of \mathbf{i} , \mathbf{j} and \mathbf{k} . [3]

.....

.....

.....

.....

.....

- (b) Find a vector equation for the line through M and N . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Find the position vector of P , the foot of the perpendicular from D to the line through M and N . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x , up to and including the term in x^2 . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

10 (a) The complex numbers v and w satisfy the equations

$$v + iw = 5 \quad \text{and} \quad (1 + 2i)v - w = 3i.$$

Solve the equations for v and w , giving your answers in the form $x + iy$, where x and y are real.

[6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) (i) On an Argand diagram, sketch the locus of points representing complex numbers z satisfying $|z - 2 - 3i| = 1$. [2]

- (ii) Calculate the least value of $\arg z$ for points on this locus. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



MATHEMATICS

9709/31

Paper 3 Pure Mathematics 3

May/June 2020

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Blank pages are indicated.

2 (a) Expand $(2 - 3x)^{-2}$ in ascending powers of x , up to and including the term in x^2 , simplifying the coefficients. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) State the set of values of x for which the expansion is valid. [1]

.....

.....

.....

.....

.....

- 3 Express the equation $\tan(\theta + 60^\circ) = 2 + \tan(60^\circ - \theta)$ as a quadratic equation in $\tan \theta$, and hence solve the equation for $0^\circ \leq \theta \leq 180^\circ$. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 The curve with equation $y = e^{2x}(\sin x + 3 \cos x)$ has a stationary point in the interval $0 \leq x \leq \pi$.

(a) Find the x -coordinate of this point, giving your answer correct to 2 decimal places. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Determine whether the stationary point is a maximum or a minimum. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5 (a) Find the quotient and remainder when $2x^3 - x^2 + 6x + 3$ is divided by $x^2 + 3$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Using your answer to part (a), find the exact value of $\int_1^3 \frac{2x^3 - x^2 + 6x + 3}{x^2 + 3} dx$. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

6



The diagram shows a circle with centre O and radius r . The tangents to the circle at the points A and B meet at T , and angle AOB is $2x$ radians. The shaded region is bounded by the tangents AT and BT , and by the minor arc AB . The area of the shaded region is equal to the area of the circle.

(a) Show that x satisfies the equation $\tan x = \pi + x$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) This equation has one root in the interval $0 < x < \frac{1}{2}\pi$. Verify by calculation that this root lies between 1 and 1.4. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Use the iterative formula

$$x_{n+1} = \tan^{-1}(\pi + x_n)$$

to determine the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

7 Let $f(x) = \frac{\cos x}{1 + \sin x}$.

(a) Show that $f'(x) < 0$ for all x in the interval $-\frac{1}{2}\pi < x < \frac{3}{2}\pi$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find $\int_{\frac{1}{6}\pi}^{\frac{1}{2}\pi} f(x) dx$. Give your answer in a simplified exact form. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

8 A certain curve is such that its gradient at a point (x, y) is proportional to $\frac{y}{x\sqrt{x}}$. The curve passes through the points with coordinates $(1, 1)$ and $(4, e)$.

(a) By setting up and solving a differential equation, find the equation of the curve, expressing y in terms of x . [8]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Describe what happens to y as x tends to infinity.

[1]

.....

.....

.....

.....

.....

- 9 With respect to the origin O , the vertices of a triangle ABC have position vectors

$$\vec{OA} = 2\mathbf{i} + 5\mathbf{k}, \quad \vec{OB} = 3\mathbf{i} + 2\mathbf{j} + 3\mathbf{k} \quad \text{and} \quad \vec{OC} = \mathbf{i} + \mathbf{j} + \mathbf{k}.$$

- (a) Using a scalar product, show that angle ABC is a right angle. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Show that triangle ABC is isosceles. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

10 (a) The complex number u is defined by $u = \frac{3i}{a + 2i}$, where a is real.

(i) Express u in the Cartesian form $x + iy$, where x and y are in terms of a . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the exact value of a for which $\arg u^* = \frac{1}{3}\pi$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) (i) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - 2i| \leq |z - 1 - i|$ and $|z - 2 - i| \leq 2$. [4]

- (ii) Calculate the least value of $\arg z$ for points in this region. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/32

Paper 3 Pure Mathematics 3

May/June 2020

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Blank pages are indicated.



- 1 Find the quotient and remainder when $6x^4 + x^3 - x^2 + 5x - 6$ is divided by $2x^2 - x + 1$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

2



The variables x and y satisfy the equation $y^2 = Ae^{kx}$, where A and k are constants. The graph of $\ln y$ against x is a straight line passing through the points $(1.5, 1.2)$ and $(5.24, 2.7)$ as shown in the diagram.

Find the values of A and k correct to 2 decimal places. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3 Find the exact value of

$$\int_1^4 x^{\frac{3}{2}} \ln x \, dx.$$

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 A curve has equation $y = \cos x \sin 2x$.

Find the x -coordinate of the stationary point in the interval $0 < x < \frac{1}{2}\pi$, giving your answer correct to 3 significant figures. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

6



The diagram shows the curve $y = \frac{x}{1 + 3x^4}$, for $x \geq 0$, and its maximum point M .

(a) Find the x -coordinate of M , giving your answer correct to 3 decimal places. [4]

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

8 (a) Solve the equation $(1 + 2i)w + iw^* = 3 + 5i$. Give your answer in the form $x + iy$, where x and y are real. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) (i) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - 2 - 2i| \leq 1$ and $\arg(z - 4i) \geq -\frac{1}{4}\pi$. [4]

(ii) Find the least value of $\text{Im } z$ for points in this region, giving your answer in an exact form. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Use the iterative formula $p_{n+1} = \tan^{-1}\left(\frac{1}{1+p_n}\right)$ to determine the value of p correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Hence find the value of k correct to 2 decimal places. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

10 With respect to the origin O , the points A and B have position vectors given by $\vec{OA} = 6\mathbf{i} + 2\mathbf{j}$ and $\vec{OB} = 2\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$. The midpoint of OA is M . The point N lying on AB , between A and B , is such that $AN = 2NB$.

(a) Find a vector equation for the line through M and N . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The line through M and N intersects the line through O and B at the point P .

(b) Find the position vector of P . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Calculate angle OPM , giving your answer in degrees. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

* 1 8 6 2 6 0 5 1 6 9 *

MATHEMATICS**9709/33**

Paper 3 Pure Mathematics 3

May/June 2020**1 hour 50 minutes**

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Blank pages are indicated.

1 Solve the inequality $|2x - 1| > 3|x + 2|$.

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

2 Find the exact value of $\int_0^1 (2-x)e^{-2x} dx$. [5]

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

- 3 (a) Show that the equation

$$\ln(1 + e^{-x}) + 2x = 0$$

can be expressed as a quadratic equation in e^x . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Hence solve the equation $\ln(1 + e^{-x}) + 2x = 0$, giving your answer correct to 3 decimal places. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 The equation of a curve is $y = x \tan^{-1}\left(\frac{1}{2}x\right)$.

(a) Find $\frac{dy}{dx}$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) The tangent to the curve at the point where $x = 2$ meets the y -axis at the point with coordinates $(0, p)$.

Find p . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5 By first expressing the equation

$$\tan \theta \tan(\theta + 45^\circ) = 2 \cot 2\theta$$

as a quadratic equation in $\tan \theta$, solve the equation for $0^\circ < \theta < 90^\circ$.

[6]

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

A series of horizontal dotted lines for writing, consisting of 28 lines spaced evenly down the page.

- 6 (a) By sketching a suitable pair of graphs, show that the equation $x^5 = 2 + x$ has exactly one real root. [2]

.....

.....

- (b) Show that if a sequence of values given by the iterative formula

$$x_{n+1} = \frac{4x_n^5 + 2}{5x_n^4 - 1}$$

converges, then it converges to the root of the equation in part (a). [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use the iterative formula with initial value $x_1 = 1.5$ to calculate the root correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

7 Let $f(x) = \frac{2}{(2x-1)(2x+1)}$.

(a) Express $f(x)$ in partial fractions.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Using your answer to part (a), show that

$$(f(x))^2 = \frac{1}{(2x-1)^2} - \frac{1}{2x-1} + \frac{1}{2x+1} + \frac{1}{(2x+1)^2}.$$

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Hence show that $\int_1^2 (f(x))^2 dx = \frac{2}{5} + \frac{1}{2} \ln\left(\frac{5}{9}\right)$. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

8 Relative to the origin O , the points A , B and D have position vectors given by

$$\vec{OA} = \mathbf{i} + 2\mathbf{j} + \mathbf{k}, \quad \vec{OB} = 2\mathbf{i} + 5\mathbf{j} + 3\mathbf{k} \quad \text{and} \quad \vec{OD} = 3\mathbf{i} + 2\mathbf{k}.$$

A fourth point C is such that $ABCD$ is a parallelogram.

(a) Find the position vector of C and verify that the parallelogram is not a rhombus. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find angle BAD , giving your answer in degrees. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Find the area of the parallelogram correct to 3 significant figures. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

9 (a) The complex numbers u and w are such that

$$u - w = 2i \quad \text{and} \quad uw = 6.$$

Find u and w , giving your answers in the form $x + iy$, where x and y are real and exact. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

- (b) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities

$$|z - 2 - 2i| \leq 2, \quad 0 \leq \arg z \leq \frac{1}{4}\pi \quad \text{and} \quad \operatorname{Re} z \leq 3. \quad [5]$$

10



A tank containing water is in the form of a hemisphere. The axis is vertical, the lowest point is A and the radius is r , as shown in the diagram. The depth of water at time t is h . At time $t = 0$ the tank is full and the depth of the water is r . At this instant a tap at A is opened and water begins to flow out at a rate proportional to \sqrt{h} . The tank becomes empty at time $t = 14$.

The volume of water in the tank is V when the depth is h . It is given that $V = \frac{1}{3}\pi(3rh^2 - h^3)$.

(a) Show that h and t satisfy a differential equation of the form

$$\frac{dh}{dt} = -\frac{B}{2rh^{\frac{1}{2}} - h^{\frac{3}{2}}},$$

where B is a positive constant.

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

* 5 9 0 7 0 8 9 7 0 6 *



MATHEMATICS

9709/31

Paper 3 Pure Mathematics 3

October/November 2020

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Blank pages are indicated.

BLANK PAGE

- 2 On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z| \geq 2$ and $|z - 1 + i| \leq 1$. [4]

3 The parametric equations of a curve are

$$x = 3 - \cos 2\theta, \quad y = 2\theta + \sin 2\theta,$$

for $0 < \theta < \frac{1}{2}\pi$.

Show that $\frac{dy}{dx} = \cot \theta$.

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 Solve the equation

$$\log_{10}(2x + 1) = 2 \log_{10}(x + 1) - 1.$$

Give your answers correct to 3 decimal places.

[6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 5 (a) By sketching a suitable pair of graphs, show that the equation $\operatorname{cosec} x = 1 + e^{-\frac{1}{2}x}$ has exactly two roots in the interval $0 < x < \pi$. [2]

.....
.....

- (b) The sequence of values given by the iterative formula

$$x_{n+1} = \pi - \sin^{-1}\left(\frac{1}{e^{-\frac{1}{2}x_n} + 1}\right),$$

with initial value $x_1 = 2$, converges to one of these roots.

Use the formula to determine this root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

8 The coordinates (x, y) of a general point of a curve satisfy the differential equation

$$x \frac{dy}{dx} = (1 - 2x^2)y,$$

for $x > 0$. It is given that $y = 1$ when $x = 1$.

Solve the differential equation, obtaining an expression for y in terms of x . [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

9 Let $f(x) = \frac{8 + 5x + 12x^2}{(1 - x)(2 + 3x)^2}$.

(a) Express $f(x)$ in partial fractions.

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x , up to and including the term in x^2 . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

10



The diagram shows the curve $y = (2 - x)e^{-\frac{1}{2}x}$, and its minimum point M .

(a) Find the exact coordinates of M . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Find the area of the shaded region bounded by the curve and the axes. Give your answer in terms of e . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



MATHEMATICS

9709/32

Paper 3 Pure Mathematics 3

October/November 2020

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Blank pages are indicated.

BLANK PAGE

1 Solve the equation

$$\ln(1 + e^{-3x}) = 2.$$

Give the answer correct to 3 decimal places.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 2 (a) Expand $\sqrt[3]{1+6x}$ in ascending powers of x , up to and including the term in x^3 , simplifying the coefficients. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) State the set of values of x for which the expansion is valid. [1]

.....

.....

.....

.....

.....

3 The variables x and y satisfy the relation $2^y = 3^{1-2x}$.

- (a) By taking logarithms, show that the graph of y against x is a straight line. State the exact value of the gradient of this line. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Find the exact x -coordinate of the point of intersection of this line with the line $y = 3x$. Give your answer in the form $\frac{\ln a}{\ln b}$, where a and b are integers. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 4 (a) Show that the equation $\tan(\theta + 60^\circ) = 2 \cot \theta$ can be written in the form

$$\tan^2 \theta + 3\sqrt{3} \tan \theta - 2 = 0. \quad [3]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



The diagram shows the curve with parametric equations

$$x = \tan \theta, \quad y = \cos^2 \theta,$$

for $-\frac{1}{2}\pi < \theta < \frac{1}{2}\pi$.

- (a) Show that the gradient of the curve at the point with parameter θ is $-2 \sin \theta \cos^3 \theta$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The gradient of the curve has its maximum value at the point P .

(b) Find the exact value of the x -coordinate of P .

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

6 The complex number u is defined by

$$u = \frac{7+i}{1-i}.$$

(a) Express u in the form $x + iy$, where x and y are real. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Show on a sketch of an Argand diagram the points A , B and C representing u , $7 + i$ and $1 - i$ respectively. [2]

(c) By considering the arguments of $7 + i$ and $1 - i$, show that

$$\tan^{-1}\left(\frac{4}{3}\right) = \tan^{-1}\left(\frac{1}{7}\right) + \frac{1}{4}\pi. \quad [3]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) State what happens to the value of x when t tends to infinity. [1]

.....

.....

.....

.....

.....

.....

.....

8 With respect to the origin O , the position vectors of the points A , B , C and D are given by

$$\vec{OA} = \begin{pmatrix} 2 \\ 1 \\ 5 \end{pmatrix}, \quad \vec{OB} = \begin{pmatrix} 4 \\ -1 \\ 1 \end{pmatrix}, \quad \vec{OC} = \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} \quad \text{and} \quad \vec{OD} = \begin{pmatrix} 3 \\ 2 \\ 3 \end{pmatrix}.$$

(a) Show that $AB = 2CD$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the angle between the directions of \vec{AB} and \vec{CD} . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence find the exact value of $\int_0^2 f(x) dx$.

[6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

10



The diagram shows the curve $y = \sqrt{x} \cos x$, for $0 \leq x \leq \frac{3}{2}\pi$, and its minimum point M , where $x = a$. The shaded region between the curve and the x -axis is denoted by R .

- (a) Show that a satisfies the equation $\tan a = \frac{1}{2a}$. [3]

.....

.....

.....

.....

.....

.....

.....

- (b) The sequence of values given by the iterative formula $a_{n+1} = \pi + \tan^{-1}\left(\frac{1}{2a_n}\right)$, with initial value $x_1 = 3$, converges to a .

Use this formula to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



MATHEMATICS

9709/33

Paper 3 Pure Mathematics 3

October/November 2020

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Blank pages are indicated.

BLANK PAGE

1 Solve the inequality $2 - 5x > 2|x - 3|$.

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 2 On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z| \geq 2$ and $|z - 1 + i| \leq 1$. [4]

3 The parametric equations of a curve are

$$x = 3 - \cos 2\theta, \quad y = 2\theta + \sin 2\theta,$$

for $0 < \theta < \frac{1}{2}\pi$.

Show that $\frac{dy}{dx} = \cot \theta$.

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 5 (a) By sketching a suitable pair of graphs, show that the equation $\operatorname{cosec} x = 1 + e^{-\frac{1}{2}x}$ has exactly two roots in the interval $0 < x < \pi$. [2]

.....
.....

- (b) The sequence of values given by the iterative formula

$$x_{n+1} = \pi - \sin^{-1} \left(\frac{1}{e^{-\frac{1}{2}x_n} + 1} \right),$$

with initial value $x_1 = 2$, converges to one of these roots.

Use the formula to determine this root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

- 6 (a) Express $\sqrt{6} \cos \theta + 3 \sin \theta$ in the form $R \cos(\theta - \alpha)$, where $R > 0$ and $0^\circ < \alpha < 90^\circ$. State the exact value of R and give α correct to 2 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Hence solve the equation $\sqrt{6} \cos \frac{1}{3}x + 3 \sin \frac{1}{3}x = 2.5$, for $0^\circ < x < 360^\circ$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

7 (a) Verify that $-1 + \sqrt{5}i$ is a root of the equation $2x^3 + x^2 + 6x - 18 = 0$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Dotted lines for writing.

9 Let $f(x) = \frac{8 + 5x + 12x^2}{(1 - x)(2 + 3x)^2}$.

(a) Express $f(x)$ in partial fractions.

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x , up to and including the term in x^2 . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

10



The diagram shows the curve $y = (2 - x)e^{-\frac{1}{2}x}$, and its minimum point M .

- (a) Find the exact coordinates of M . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the area of the shaded region bounded by the curve and the axes. Give your answer in terms of e . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

11 Two lines have equations $\mathbf{r} = \mathbf{i} + 2\mathbf{j} + \mathbf{k} + \lambda(a\mathbf{i} + 2\mathbf{j} - \mathbf{k})$ and $\mathbf{r} = 2\mathbf{i} + \mathbf{j} - \mathbf{k} + \mu(2\mathbf{i} - \mathbf{j} + \mathbf{k})$, where a is a constant.

(a) Given that the two lines intersect, find the value of a and the position vector of the point of intersection. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



MATHEMATICS

9709/32

Paper 3 Pure Mathematics 3

February/March 2021

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Any blank pages are indicated.

BLANK PAGE

- 2 The polynomial $ax^3 + 5x^2 - 4x + b$, where a and b are constants, is denoted by $p(x)$. It is given that $(x + 2)$ is a factor of $p(x)$ and that when $p(x)$ is divided by $(x + 1)$ the remainder is 2.

Find the values of a and b .

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 3 By first expressing the equation $\tan(x + 45^\circ) = 2 \cot x + 1$ as a quadratic equation in $\tan x$, solve the equation for $0^\circ < x < 180^\circ$. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 4 The variables x and y satisfy the differential equation

$$(1 - \cos x) \frac{dy}{dx} = y \sin x.$$

It is given that $y = 4$ when $x = \pi$.

- (a) Solve the differential equation, obtaining an expression for y in terms of x . [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Sketch the graph of y against x for $0 < x < 2\pi$.

[1]

(b) Hence solve the equation $\sqrt{7} \sin 2\theta + 2 \cos 2\theta = 1$, for $0^\circ < \theta < 180^\circ$. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

6 Let $f(x) = \frac{5a}{(2x - a)(3a - x)}$, where a is a positive constant.

(a) Express $f(x)$ in partial fractions.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence show that $\int_a^{2a} f(x) \, dx = \ln 6$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

7 Two lines have equations $\mathbf{r} = \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix} + s \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix}$ and $\mathbf{r} = \begin{pmatrix} 2 \\ 1 \\ 4 \end{pmatrix} + t \begin{pmatrix} 1 \\ -1 \\ 4 \end{pmatrix}$.

(a) Show that the lines are skew.

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

8 The complex numbers u and v are defined by $u = -4 + 2i$ and $v = 3 + i$.

(a) Find $\frac{u}{v}$ in the form $x + iy$, where x and y are real. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence express $\frac{u}{v}$ in the form $re^{i\theta}$, where r and θ are exact. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

In an Argand diagram, with origin O , the points A , B and C represent the complex numbers u , v and $2u + v$ respectively.

(c) State fully the geometrical relationship between OA and BC . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) Prove that angle $AOB = \frac{3}{4}\pi$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

9 Let $f(x) = \frac{e^{2x} + 1}{e^{2x} - 1}$, for $x > 0$.

(a) The equation $x = f(x)$ has one root, denoted by a .

Verify by calculation that a lies between 1 and 1.5. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Use an iterative formula based on the equation in part (a) to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

10



The diagram shows the curve $y = \sin 2x \cos^2 x$ for $0 \leq x \leq \frac{1}{2}\pi$, and its maximum point M .

- (a) Using the substitution $u = \sin x$, find the exact area of the region bounded by the curve and the x -axis. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



MATHEMATICS

9709/31

Paper 3 Pure Mathematics 3

May/June 2021

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Any blank pages are indicated.

1 Solve the inequality $2|3x - 1| < |x + 1|$.

[4]

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

- 2 Find the real root of the equation $\frac{2e^x + e^{-x}}{2 + e^x} = 3$, giving your answer correct to 3 decimal places. Your working should show clearly that the equation has only one real root. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 3 (a) Given that $\cos(x - 30^\circ) = 2 \sin(x + 30^\circ)$, show that $\tan x = \frac{2 - \sqrt{3}}{1 - 2\sqrt{3}}$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Hence solve the equation

$$\cos(x - 30^\circ) = 2 \sin(x + 30^\circ),$$

for $0^\circ < x < 360^\circ$.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 4 (a) Prove that $\frac{1 - \cos 2\theta}{1 + \cos 2\theta} \equiv \tan^2 \theta$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Hence find the exact value of $\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} \frac{1 - \cos 2\theta}{1 + \cos 2\theta} d\theta$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5 (a) Solve the equation $z^2 - 2piz - q = 0$, where p and q are real constants. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

In an Argand diagram with origin O , the roots of this equation are represented by the distinct points A and B .

(b) Given that A and B lie on the imaginary axis, find a relation between p and q . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

6 The parametric equations of a curve are

$$x = \ln(2 + 3t), \quad y = \frac{t}{2 + 3t}.$$

(a) Show that the gradient of the curve is always positive. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

7



The diagram shows the curve $y = \frac{\tan^{-1} x}{\sqrt{x}}$ and its maximum point M where $x = a$.

(a) Show that a satisfies the equation

$$a = \tan\left(\frac{2a}{1+a^2}\right). \quad [4]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Verify by calculation that a lies between 1.3 and 1.5. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use an iterative formula based on the equation in part (a) to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 8 With respect to the origin O , the points A and B have position vectors given by $\overrightarrow{OA} = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$ and $\overrightarrow{OB} = \begin{pmatrix} 3 \\ 1 \\ -2 \end{pmatrix}$. The line l has equation $\mathbf{r} = \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix}$.

(a) Find the acute angle between the directions of AB and l . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the position vector of the point P on l such that $AP = BP$.

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

9 The equation of a curve is $y = x^{-\frac{2}{3}} \ln x$ for $x > 0$. The curve has one stationary point.

(a) Find the exact coordinates of the stationary point. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Show that $\int_1^8 y \, dx = 18 \ln 2 - 9.$

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

10 The variables x and t satisfy the differential equation $\frac{dx}{dt} = x^2(1 + 2x)$, and $x = 1$ when $t = 0$.

Using partial fractions, solve the differential equation, obtaining an expression for t in terms of x .

[11]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

A series of horizontal dotted lines for writing, consisting of 20 rows.



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



MATHEMATICS

9709/32

Paper 3 Pure Mathematics 3

May/June 2021

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Any blank pages are indicated.

1 Solve the inequality $|2x - 1| < 3|x + 1|$.

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 2 On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z + 1 - i| \leq 1$ and $\arg(z - 1) \leq \frac{3}{4}\pi$. [4]

3 The variables x and y satisfy the equation $x = A(3^{-y})$, where A is a constant.

(a) Explain why the graph of y against $\ln x$ is a straight line and state the exact value of the gradient of the line. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

It is given that the line intersects the y -axis at the point where $y = 1.3$.

(b) Calculate the value of A , giving your answer correct to 2 decimal places. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 Using integration by parts, find the exact value of $\int_0^2 \tan^{-1}\left(\frac{1}{2}x\right) dx$. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

6 (a) Prove that $\operatorname{cosec} 2\theta - \cot 2\theta \equiv \tan \theta$.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence show that $\int_{\frac{1}{4}\pi}^{\frac{1}{3}\pi} (\operatorname{cosec} 2\theta - \cot 2\theta) \, d\theta = \frac{1}{2} \ln 2$.

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

A series of horizontal dotted lines for writing, consisting of 25 lines spaced evenly down the page.

- 8 The equation of a curve is $y = e^{-5x} \tan^2 x$ for $-\frac{1}{2}\pi < x < \frac{1}{2}\pi$.

Find the x -coordinates of the stationary points of the curve. Give your answers correct to 3 decimal places where appropriate. [8]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Ruled lines for writing.

9 Let $f(x) = \frac{14 - 3x + 2x^2}{(2 + x)(3 + x^2)}$.

(a) Express $f(x)$ in partial fractions.

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

10



The diagram shows a trapezium $ABCD$ in which $AD = BC = r$ and $AB = 2r$. The acute angles BAD and ABC are both equal to x radians. Circular arcs of radius r with centres A and B meet at M , the midpoint of AB .

- (a) Given that the sum of the areas of the shaded sectors is 90% of the area of the trapezium, show that x satisfies the equation $x = 0.9(2 - \cos x) \sin x$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Verify by calculation that x lies between 0.5 and 0.7. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Show that if a sequence of values in the interval $0 < x < \frac{1}{2}\pi$ given by the iterative formula

$$x_{n+1} = \cos^{-1} \left(2 - \frac{x_n}{0.9 \sin x_n} \right)$$

converges, then it converges to the root of the equation in part (a). [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) Use this iterative formula to determine x correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



MATHEMATICS

9709/43

Paper 4 Mechanics

May/June 2021

1 hour 15 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity (g) is needed, use 10 m s^{-2} .

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **12** pages.

- 1 Particles P of mass 0.4 kg and Q of mass 0.5 kg are free to move on a smooth horizontal plane. P and Q are moving directly towards each other with speeds 2.5 m s^{-1} and 1.5 m s^{-1} respectively. After P and Q collide, the speed of Q is twice the speed of P .

Find the two possible values of the speed of P after the collision.

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

2 A cyclist is travelling along a straight horizontal road. She is working at a constant rate of 150 W. At an instant when her speed is 4 m s^{-1} , her acceleration is 0.25 m s^{-2} . The resistance to motion is 20 N.

(a) Find the total mass of the cyclist and her bicycle. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The cyclist comes to a straight hill inclined at an angle θ above the horizontal. She ascends the hill at constant speed 3 m s^{-1} . She continues to work at the same rate as before and the resistance force is unchanged.

(b) Find the value of θ . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

3



Four coplanar forces act at a point. The magnitudes of the forces are 20 N, 30 N, 40 N and F N. The directions of the forces are as shown in the diagram, where $\sin \alpha^\circ = 0.28$ and $\sin \beta^\circ = 0.6$.

Given that the forces are in equilibrium, find F and θ .

[6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 A particle is projected vertically upwards with speed $u \text{ m s}^{-1}$ from a point on horizontal ground. After 2 seconds, the height of the particle above the ground is 24 m.

(a) Show that $u = 22$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) The height of the particle above the ground is more than h m for a period of 3.6 s.
Find h . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5 A car of mass 1400 kg is towing a trailer of mass 500 kg down a straight hill inclined at an angle of 5° to the horizontal. The car and trailer are connected by a light rigid tow-bar. At the top of the hill the speed of the car and trailer is 20 m s^{-1} and at the bottom of the hill their speed is 30 m s^{-1} .

(a) It is given that as the car and trailer descend the hill, the engine of the car does 150 000 J of work, and there are no resistance forces.

Find the length of the hill. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 6 A particle moves in a straight line and passes through the point A at time $t = 0$. The velocity of the particle at time t s after leaving A is v m s⁻¹, where

$$v = 2t^2 - 5t + 3.$$

- (a) Find the times at which the particle is instantaneously at rest. Hence or otherwise find the minimum velocity of the particle. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Sketch the velocity-time graph for the first 3 seconds of motion. [3]

(c) Find the distance travelled between the two times when the particle is instantaneously at rest. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



A particle P of mass 0.3 kg rests on a rough plane inclined at an angle θ to the horizontal, where $\sin \theta = \frac{7}{25}$. A horizontal force of magnitude 4 N , acting in the vertical plane containing a line of greatest slope of the plane, is applied to P (see diagram). The particle is on the point of sliding up the plane.

- (a) Show that the coefficient of friction between the particle and the plane is $\frac{3}{4}$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The force acting horizontally is replaced by a force of magnitude 4 N acting up the plane parallel to a line of greatest slope.

- (b) Find the acceleration of P . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Starting with P at rest, the force of 4 N parallel to the plane acts for 3 seconds and is then removed.

Find the total distance travelled until P comes to instantaneous rest. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

* 0 7 4 6 8 9 6 0 5 1 *



MATHEMATICS

9709/31

Paper 3 Pure Mathematics 3

October/November 2021

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages.

1 Solve the equation $4|5^x - 1| = 5^x$, giving your answers correct to 3 decimal places. [4]

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

- 2 (a) Express $5 \sin x - 3 \cos x$ in the form $R \sin(x - \alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$. Give the exact value of R and give α correct to 2 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Hence state the greatest and least possible values of $(5 \sin x - 3 \cos x)^2$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3 The curve with equation $y = xe^{1-2x}$ has one stationary point.

(a) Find the coordinates of this point. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Determine whether the stationary point is a maximum or a minimum. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 Using the substitution $u = \sqrt{x}$, find the exact value of

$$\int_3^{\infty} \frac{1}{(x+1)\sqrt{x}} dx. \quad [6]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5 (a) Show that the equation

$$\cot 2\theta + \cot \theta = 2$$

can be expressed as a quadratic equation in $\tan \theta$.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence solve the equation $\cot 2\theta + \cot \theta = 2$, for $0 < \theta < \pi$, giving your answers correct to 3 decimal places.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 6 When $(a + bx)\sqrt{1 + 4x}$, where a and b are constants, is expanded in ascending powers of x , the coefficients of x and x^2 are 3 and -6 respectively.

Find the values of a and b .

[6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

7 (a) Given that $y = \ln(\ln x)$, show that

$$\frac{dy}{dx} = \frac{1}{x \ln x}. \quad [1]$$

.....

.....

.....

.....

.....

The variables x and t satisfy the differential equation

$$x \ln x + t \frac{dx}{dt} = 0.$$

It is given that $x = e$ when $t = 2$.

(b) Solve the differential equation obtaining an expression for x in terms of t , simplifying your answer. [7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Hence state what happens to the value of x as t tends to infinity. [1]

.....

.....

.....

.....

(b) Verify by calculation that a lies between 9 and 11. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use an iterative formula based on the equation in part (a) to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

9 Two lines l and m have equations $\mathbf{r} = 3\mathbf{i} + 2\mathbf{j} + 5\mathbf{k} + s(4\mathbf{i} - \mathbf{j} + 3\mathbf{k})$ and $\mathbf{r} = \mathbf{i} - \mathbf{j} - 2\mathbf{k} + t(-\mathbf{i} + 2\mathbf{j} + 2\mathbf{k})$ respectively.

(a) Show that l and m are perpendicular. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Show that l and m intersect and state the position vector of the point of intersection. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Show that the length of the perpendicular from the origin to the line m is $\frac{1}{3}\sqrt{5}$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

10 The complex number $1 + 2i$ is denoted by u . The polynomial $2x^3 + ax^2 + 4x + b$, where a and b are real constants, is denoted by $p(x)$. It is given that u is a root of the equation $p(x) = 0$.

(a) Find the values of a and b . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) State a second complex root of this equation. [1]

.....

.....

.....

.....

(c) Find the real factors of $p(x)$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) (i) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - u| \leq \sqrt{5}$ and $\arg z \leq \frac{1}{4}\pi$. [4]

(ii) Find the least value of $\text{Im } z$ for points in the shaded region. Give your answer in an exact form. [1]

.....

.....

.....

.....

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



MATHEMATICS

9709/32

Paper 3 Pure Mathematics 3

October/November 2021

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Any blank pages are indicated.

BLANK PAGE

- 1 Find the value of x for which $3(2^{1-x}) = 7^x$. Give your answer in the form $\frac{\ln a}{\ln b}$, where a and b are integers. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

2 Solve the inequality $|3x - a| > 2|x + 2a|$, where a is a positive constant.

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 3 (a) Given the complex numbers $u = a + ib$ and $w = c + id$, where a , b , c and d are real, prove that $(u + w)^* = u^* + w^*$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Solve the equation $(z + 2 + i)^* + (2 + i)z = 0$, giving your answer in the form $x + iy$ where x and y are real. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 Express $\frac{4x^2 - 13x + 13}{(2x - 1)(x - 3)}$ in partial fractions.

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 5 (a) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - 3 - 2i| \leq 1$ and $\text{Im } z \geq 2$. [4]

- (b) Find the greatest value of $\arg z$ for points in the shaded region, giving your answer in degrees. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

6 (a) Using the expansions of $\sin(3x + 2x)$ and $\sin(3x - 2x)$, show that

$$\frac{1}{2}(\sin 5x + \sin x) \equiv \sin 3x \cos 2x. \qquad [3]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence show that $\int_0^{\frac{1}{4}\pi} \sin 3x \cos 2x \, dx = \frac{1}{5}(3 - \sqrt{2})$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

7 The variables x and y satisfy the differential equation

$$e^{2x} \frac{dy}{dx} = 4xy^2,$$

and it is given that $y = 1$ when $x = 0$.

Solve the differential equation, obtaining an expression for y in terms of x .

[7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the exact coordinates of the point on the curve where the tangent is parallel to the y-axis.

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

10 With respect to the origin O , the position vectors of the points A and B are given by $\vec{OA} = \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}$ and $\vec{OB} = \begin{pmatrix} 0 \\ 3 \\ 1 \end{pmatrix}$.

(a) Find a vector equation for the line l through A and B . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) The point C lies on l and is such that $\vec{AC} = 3\vec{AB}$.
Find the position vector of C . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

11 The equation of a curve is $y = \sqrt{\tan x}$, for $0 \leq x < \frac{1}{2}\pi$.

(a) Express $\frac{dy}{dx}$ in terms of $\tan x$, and verify that $\frac{dy}{dx} = 1$ when $x = \frac{1}{4}\pi$. [4]

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

The value of $\frac{dy}{dx}$ is also 1 at another point on the curve where $x = a$, as shown in the diagram.



(b) Show that $t^3 + t^2 + 3t - 1 = 0$, where $t = \tan a$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use the iterative formula

$$a_{n+1} = \tan^{-1} \left(\frac{1}{3} (1 - \tan^2 a_n - \tan^3 a_n) \right)$$

to determine a correct to 2 decimal places, giving the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



MATHEMATICS

9709/33

Paper 3 Pure Mathematics 3

October/November 2021

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Any blank pages are indicated.

2 (a) Sketch the graph of $y = |2x - 3|$.

[1]

(b) Solve the inequality $|2x - 3| < 3x + 2$.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

6 (a) By first expanding $\cos(x - 60^\circ)$, show that the expression

$$2 \cos(x - 60^\circ) + \cos x$$

can be written in the form $R \cos(x - \alpha)$, where $R > 0$ and $0^\circ < \alpha < 90^\circ$. Give the exact value of R and the value of α correct to 2 decimal places. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence find the value of x in the interval $0^\circ < x < 360^\circ$ for which $2 \cos(x - 60^\circ) + \cos x$ takes its least possible value. [2]

.....

.....

.....

.....

.....

7 The equation of a curve is $\ln(x + y) = x - 2y$.

(a) Show that $\frac{dy}{dx} = \frac{x + y - 1}{2(x + y) + 1}$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



In the diagram, $OABCD$ is a pyramid with vertex D . The horizontal base $OABC$ is a square of side 4 units. The edge OD is vertical and $OD = 4$ units. The unit vectors \mathbf{i} , \mathbf{j} and \mathbf{k} are parallel to OA , OC and OD respectively.

The midpoint of AB is M and the point N on CD is such that $DN = 3NC$.

- (a) Find a vector equation for the line through M and N . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Using the substitution $u = \sqrt{x}$, show that $\int_0^4 f(x) dx = \frac{1}{3} \ln 5$. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 10 A large plantation of area 20 km^2 is becoming infected with a plant disease. At time t years the area infected is $x \text{ km}^2$ and the rate of increase of x is proportional to the ratio of the area infected to the area not yet infected.

When $t = 0$, $x = 1$ and $\frac{dx}{dt} = 1$.

- (a) Show that x and t satisfy the differential equation

$$\frac{dx}{dt} = \frac{19x}{20 - x}. \quad [2]$$

.....

.....

.....

.....

.....

.....

.....

- (b) Solve the differential equation and show that when $t = 1$ the value of x satisfies the equation $x = e^{0.9+0.05x}$. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use an iterative formula based on the equation in part (b), with an initial value of 2, to determine x correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) Calculate the value of t at which the entire plantation becomes infected. [1]

.....

.....

.....

.....

11 The complex number $-\sqrt{3} + i$ is denoted by u .

(a) Express u in the form $re^{i\theta}$, where $r > 0$ and $-\pi < \theta \leq \pi$, giving the exact values of r and θ . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence show that u^6 is real and state its value. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) (i) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $0 \leq \arg(z - u) \leq \frac{1}{4}\pi$ and $\operatorname{Re} z \leq 2$. [4]

- (ii) Find the greatest value of $|z|$ for points in the shaded region. Give your answer correct to 3 significant figures. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

